

CURRENT PROTOCOLS IN MOLECULAR BIOLOGY

VOLUME 1

EDITORIAL BOARD

Frederick M. Ausubel
Massachusetts General Hospital & Harvard Medical School

Roger Brent
Massachusetts General Hospital & Harvard Medical School

Robert E. Kingston
Massachusetts General Hospital & Harvard Medical School

David D. Moore
Massachusetts General Hospital & Harvard Medical School

J.G. Seidman
Harvard Medical School

John A. Smith
University of Alabama

Kevin Struhl
Harvard Medical School

GUEST EDITORS

Lisa M. Albright
DNA Sequencing

Donald M. Coen
Harvard Medical School
Polymerase Chain Reaction

Ajit Varki
University of California San Diego
Glycoproteins

SERIES EDITOR

Virginia Benson Chanda



John Wiley & Sons, Inc.

CORE 13 (S36)

Copyright © 1994–1997 by John Wiley & Sons, Inc.

Copyright © 1987–1994 by Current Protocols

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

While the authors, editors, and publisher believe that the specification and usage of reagents, equipment, and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of experimental reagents, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each chemical, piece of equipment, reagent, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. This is particularly important in regard to new or infrequently employed chemicals or experimental reagents.

Library of Congress Cataloging in Publication Data:

Current protocols in molecular biology. 3 vols.

1. Molecular biology—Technique. 2. Molecular biology—Laboratory manuals. I. Ausubel, Frederick M.

QH506.C87 1987 574.8'8'028 87-21033
ISBN 0-471-50338-X

Printed in the United States of America

20 19 18 17 16 15 14 13

IN SITU HYBRIDIZATION AND IMMUNOHISTOCHEMISTRY

14

INTRODUCTION	14.0.3
14.1 Fixation, Embedding, and Sectioning of Tissues, Embryos, and Single Cells	14.1.1
Basic Protocol: Paraformaldehyde Fixation and Paraffin Wax Embedding of Tissues and Embryos	14.1.1
Alternate Protocol: Fixation of Suspended and Cultured Cells	14.1.3
Support Protocol: Perfusion of Adult Mice	14.1.4
Support Protocol: Sectioning Samples in Wax Blocks	14.1.4
Reagents and Solutions	14.1.6
Commentary	14.1.7
14.2 Cryosectioning	14.2.1
Basic Protocol: Specimen Preparation and Sectioning	14.2.1
Support Protocol: Fixation of Cryosections for In situ Hybridization	14.2.5
Support Protocol: Tissue Fixation and Sucrose Infusion	14.2.6
Commentary	14.2.6
14.3 In situ Hybridization to Cellular RNA	14.3.1
Basic Protocol: Hybridization Using Paraffin Sections and Cells	14.3.1
Alternate Protocol: Hybridization Using Cryosections	14.3.5
Support Protocol: Synthesis of ³⁵ S-Labeled Riboprobes	14.3.7
Support Protocol: Synthesis of ³⁵ S-Labeled Double-Stranded DNA Probes	14.3.8
Reagents and Solutions	14.3.8
Commentary	14.3.11
14.4 Detection of Hybridized Probe	14.4.1
Basic Protocol: Film Autoradiography	14.4.1
Basic Protocol: Emulsion Autoradiography	14.4.1
Support Protocol: Preparation of Diluted Emulsion for Autoradiography	14.4.2
Commentary	14.4.3
14.5 Counterstaining and Mounting of Autoradiographed In situ Hybridization Slides	14.5.1
Basic Protocol: Giemsa Staining	14.5.1
Alternate Protocol: Hematoxylin/Eosin Staining	14.5.2
Alternate Protocol: Toluidine Blue Staining	14.5.3
Alternate Protocol: Hoechst Staining	14.5.3
Reagents and Solutions	14.5.4
Commentary	14.5.5
14.6 Immunohistochemistry	14.6.1
Basic Protocol: Immunofluorescent Labeling of Cells Grown as Monolayers	14.6.1
Alternate Protocol: Immunofluorescent Labeling of Suspension Cells	14.6.2
Basic Protocol: Immunofluorescent Labeling of Tissue Sections	14.6.3
Alternate Protocol: Immunofluorescent Labeling of Tissue Sections Using Coplin Jars	14.6.4
Alternate Protocol: Immunofluorescent Labeling Using Streptavidin-Biotin Conjugates	14.6.5
Alternate Protocol: Immunogold Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunoperoxidase Labeling of Tissue Sections	14.6.6
Alternate Protocol: Immunofluorescent Double-Labeling of Tissue Sections	14.6.7
Reagents and Solutions	14.6.7
Commentary	14.6.8

continued

- 1c. *Harsh treatment:* Pour several hundred milliliters of boiling 0.1% SDS onto the membrane. Cool to room temperature.

If a membrane is to be reprobbed, it must not be allowed to dry out between hybridization and stripping. If it becomes dry, the probe may bind to the matrix.

2. Place membrane on a sheet of dry Whatman 3MM filter paper and blot excess liquid with a second sheet. Wrap the membrane in plastic wrap and set up an autoradiograph.

If signal is still seen after autoradiography, rewash using harsher conditions.

3. The membrane can now be rehybridized. Alternatively, it can be dried and stored for later use.

Membranes can be stored dry between Whatman 3MM paper for several months at room temperature. For long-term storage, place the membranes in a desiccator at room temperature or 4°C.

REAGENTS AND SOLUTIONS

Aqueous prehybridization/hybridization (APH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see below) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Denatured salmon sperm DNA

Dissolve 10 mg Sigma type III salmon sperm DNA (sodium salt) in 1 ml water. Pass vigorously through a 17-G needle 20 times to shear the DNA. Place in a boiling water bath for 10 min, then chill. Use immediately or store at -20°C in small aliquots. If stored, reheat to 100°C for 5 min and chill on ice immediately before using.

Formamide prehybridization/hybridization (FPH) solution

5× SSC (APPENDIX 2)

5× Denhardt solution (APPENDIX 2)

50% (w/v) formamide

1% (w/v) SDS

Add 100 µg/ml denatured salmon sperm DNA (see above) just before use

Alternatives to Denhardt solution and denatured salmon sperm DNA as blocking agents are listed in Table 2.10.5 (see discussion in critical parameters).

Commercial formamide is usually satisfactory for use. If the liquid has a yellow color, deionize as follows: add 5 g of mixed-bed ion-exchange resin [e.g., Bio-Rad AG 501-X8 or 501-X8(D) resins] per 100 ml formamide, stir at room temperature for 1 hr, and filter through Whatman no. 1 paper.

CAUTION: Formamide is a teratogen. Handle with care.

Labeling buffer

200 mM Tris-Cl, pH 7.5

30 mM MgCl₂

10 mM spermidine

Mild stripping solution

5 mM Tris-Cl, pH 8.0

2 mM EDTA

0.1× Denhardt solution (APPENDIX 2)